



ISSN: 2339-0883

# **SEMINAR TAHUNAN HASIL PENELITIAN PERIKANAN DAN KELAUTAN VI ANNUAL SEMINAR OF FISHERIES AND MARINE SCIENCE VI**

## **PROSIDING**

**APLIKASI IPTEK PERIKANAN DAN KELAUTAN DALAM PENGELOLAAN,  
MITIGASI BENCANA DAN DEGRADASI WILAYAH PESISIR,  
LAUT DAN PULAU-PULAU KECIL**

**APPLICATION OF FISHERIES AND MARINE SCIENCE AND TECHNOLOGY  
ON MANAGEMENT, MITIGATION OF DISASTER  
AND ENVIRONMENTAL DEGRADATION  
IN COASTAL AREAS, SEAS AND SMALL ISLANDS**

**SEMARANG, 12 NOVEMBER 2016**

**FAKULTAS PERIKANAN DAN ILMU KELAUTAN  
UNIVERSITAS DIPONEGORO  
JUNI, 2017**

## KATA PENGANTAR

Tahun 2016 merupakan seminar tahunan ke VI yang diselenggarakan oleh FPIK UNDIP. Kegiatan seminar ini telah dimulai sejak tahun 2007 dan dilaksanakan secara berkala. Tema kegiatan seminar dari tahun ketahun bervariasi mengikuti perkembangan isu terkini di sektor perikanan dan kelautan.

Kegiatan seminar ini merupakan salah satu bentuk kontribusi perguruan tinggi khususnya FPIK UNDIP dalam upaya mendukung pembangunan di sektor perikanan dan kelautan. IPTEK sangat diperlukan untuk mendukung pembangunan sehingga tujuan pembangunan dapat tercapai dan bermanfaat bagi kemakmuran rakyat.

Dalam implementasi pembangunan selalu ada dampak yang ditimbulkan. Untuk itu, diperlukan suatu upaya agar dampak negatif dapat diminimalisir atau bahkan tidak terjadi. Oleh karena itu, Seminar ini bertemakan tentang **Aplikasi IPTEK Perikanan dan Kelautan dalam Mitigasi Bencana dan Degradasi Wilayah Pesisir, Laut dan Pulau-Pulau Kecil**. Pada kesempatan kali ini, diharapkan IPTEK hasil penelitian mengenai pengelolaan, mitigasi bencana dan degradasi wilayah pesisir, laut dan pulau-pulau kecil dapat terpublikasikan sehingga dapat dimanfaatkan untuk pembangunan yang berkelanjutan dan dapat menjaga kelestarian lingkungan. Seminar Tahunan Hasil Penelitian Perikanan dan Kelautan ke-VI merupakan kolaborasi FPIK UNDIP dan Pusat Kajian Mitigasi Bencana dan Rehabilitasi Pesisir (PKMBRP) UNDIP.

Pada kesempatan ini kami selaku panitia penyelenggara mengucapkan terimakasih kepada pemakalah, reviewer, peserta serta Pertamina EP Asset 3 Tambun Field yang telah mendukung kegiatan Seminar Tahunan Penelitian Hasil Penelitian Perikanan dan Kelautan VI sehingga dapat terlaksana dengan baik. Harapan kami semoga hasil seminar ini dapat memberikan kontribusi dalam upaya mitigasi bencana dan rehabilitasi pesisir, laut dan pulau-pulau kecil.

Semarang, Juni 2017

Panitia





## SUSUNAN PANITIA SEMINAR

Pembina	: Dekan FPIK Undip Prof. Dr. Ir. Agus Sabdono, M.Sc
Penanggung jawab	: Wakil Dekan Bidang IV Tita Elvita Sari, S.Pi., M.Sc., Ph.D
Ketua	: Dr.Sc. Anindya Wirasatriya, ST, M.Si., M.Sc
Wakil Ketua	: Dr.Ir. Suryanti, M.Pi
Sekretaris I	: Faik Kurohman, S.Pi, M.Si
Sekretaris II	: Wiwiet Teguh T, SPi, MSi
Bendahara I	: Ir. Nirwani, MSi
Bendahara II	: Retno Ayu K, S.Pi., M.Sc
Kesekretariatan	: 1. Dr. Agus Trianto, ST., M.Sc 2. Dr. Denny Nugroho, ST, M.Si 3. Kukuh Eko Prihantoko, S.Pi., M.Si 4. Sigit Febrianto, S.Kel., M.Si 5. Lukita P., STP, M.Sc 6. Lilik Maslukah, ST., M.Si 7. Ir. Ria Azizah, M.Si
Acara dan Sidang	: 1. Dr. Aristi Dian P.F., S.Pi., M.Si 2. Dr. Ir. Diah Permata W., M.Sc 3. Ir. Retno Hartati, M.Sc 4. Dr. Muhammad Helmi, S.Si., M.Si
Konsumsi	: 1. Ir. Siti Rudiyantri, M.Si 2. Ir. Sri Redjeki, M.Si 3. Ir. Ken Suwartimah, M.Si
Perlengkapan	: 1. Bogi Budi J., S.Pi., M.Si 2. A. Harjuno Condro, S.Pi, M.Si



**DEWAN REDAKSI**  
**PROSIDING**  
**SEMINAR NASIONAL TAHUNAN KE-VI**  
**HASIL-HASIL PENELITIAN PERIKANAN DAN KELAUTAN**

- Diterbitkan oleh : Fakultas Perikanan dan Ilmu Kelautan, Universitas Diponegoro  
bekerjasama dengan Pusat Kajian Mitigasi Bencana dan  
Rehabilitasi Pesisir serta Pertamina EP Asset 3 Tambun Field
- Penanggung jawab : Dekan FPIK Undip  
(Prof. Dr. Ir. Agus Sabdono, M.Sc)  
Wakil Dekan Bidang IV  
(Tita Elvita Sari, S.Pi., M.Sc., Ph.D)
- Pengarah : 1. Dr. Denny Nugroho, ST, M.Si (Kadept. Oceanografi)  
2. Dr. Ir. Diah Permata W., M.Sc (Kadept. Ilmu Kelautan)  
3. Dr. Ir. Haeruddin, M.Si (Kadept. Manajemen SD. Akuatik)  
4. Dr. Aristi Dian P.F., S.Pi., M.Si (Kadept. Perikanan Tangkap)  
5. Dr. Ir. Eko Nur C, M.Sc (Kadept. Teknologi Hasil Perikanan)  
6. Dr. Ir. Sardjito, M.App.Sc (Kadept. Akuakultur)
- Tim Editor : 1. Dr. Sc. Anindya Wirasatriya, ST, M.Si., M.Sc  
2. Dr. Ir. Suryanti, M.Pi  
3. Faik Kurohman, S.Pi, Msi  
4. Wiwiet Teguh T, S.Pi., M.Si  
5. Ir. Nirwani, Msi  
6. Retno Ayu K, S.Pi., M.Sc  
7. Dr. Aristi Dian P.F., S.Pi., M.Si  
8. Dr. Ir. Diah Permata W., M.Sc  
9. Ir. Retno Hartati, M.Sc  
10. Dr. Muhammad Helmi, S.Si., M.Si
- Reviewer : 1. Dr. Agus Trianto, ST., M.Sc  
2. Dr. Denny Nugroho, ST, M.Si  
3. Sigit Febrianto, S.Kel., M.Si  
4. Lukita P., STP, M.Sc  
5. Ir. Ria Azizah, M.Si  
6. Lilik Maslukah, ST., M.Si  
7. Ir. Siti Rudiyan, M.Si  
8. Ir. Sri Redjeki, M.Si  
9. Ir. Ken Suwartimah, M.Si  
10. Bogi Budi J., S.Pi., M.Si  
11. A. Harjuno Condro, S.Pi, M.Si
- Desain sampul : Kuku Eko Prihantoko, S.Pi., M.Si
- Layout dan tata letak : Divta Pratama Yudistira
- Alamat redaksi : Fakultas Perikanan dan Ilmu Kelautan  
Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275  
Telpn/ Fax: 024 7474698



## DAFTAR ISI

	halaman
HALAMAN JUDUL.....	i
KATA PENGANTAR .....	ii
SUSUNAN PANITIA SEMINAR .....	iii
DEWAN REDAKSI.....	iv
DAFTAR ISI .....	v

### Aplikasi IPTEK Perikanan dan Kelautan dalam Pengelolaan dan Pemanfaatan Sumberdaya Wilayah Pesisir, Laut dan Pulau-pulau Kecil (Pemanfaatan Sumberdaya Perairan)

1. Research About Stock Condition of Skipjack Tuna ( <i>Katsuwonus pelamis</i> ) in Gulf of Bone South Sulawesi, Indonesia .....	1
2. Keberhasilan Usaha Pemberdayaan Ekonomi Kelompok Perajin Batik Mangrove dalam Perbaikan Mutu dan Peningkatan Hasil Produksi di Mangkang Wetan, Semarang .....	15
3. Pengelolaan Perikanan Cakalang Berkelanjutan Melalui Studi Optimalisasi dan Pendekatan Bioekonomi di Kota Kendari .....	22
4. Kajian Pengembangan Desa Pantai Mekar, Kecamatan Muara Gembong, Kabupaten Bekasi sebagai Kampung Wisata Bahari .....	33
5. Kajian Valuasi Ekonomi Hutan Mangrove di Desa Pantai Mekar, Kecamatan Muara Gembong, Kabupaten Bekasi.....	47
6. Studi Pemetaan Aset Nelayan di Desa Pantai Mekar, Kecamatan Muara Gembong, Kabupaten Bekasi .....	55
7. Hubungan Antara Daerah Penangkapan Rajungan ( <i>Portunus pelagicus</i> ) dengan Parameter Oseanografi di Perairan Tegal, Jawa Tengah .....	67
8. Komposisi Jenis Hiu dan Distribusi Titik Penangkapannya di Perairan Pesisir Cilacap, Jawa Tengah.....	82
9. Analisis Pengembangan Fasilitas Pelabuhan yang Berwawasan Lingkungan ( <i>Ecoport</i> ) di Pelabuhan Perikanan Nusantara (PPN) Pengembangan, Jembrana Bali.....	93
10. Anallisis Kepuasan Pengguna Pelabuhan Perikanan Nusantara (PPN) Pengembangan, Jembrana Bali .....	110
11. Effect of Different Soaking Time in Coconut Shell Liquid Smoke to The Profile of Lipids Cats Fish ( <i>Clarias batrachus</i> ) Smoke.....	124



## Rehabilitasi Ekosistem: Mangrove, Terumbu Karang dan Padang Lamun

1. Pola Pertumbuhan, Respon Osmotik dan Tingkat Kematangan Gonad Kerang *Polymesoda erosa* di Perairan Teluk Youtefa Jayapura Papua ..... 135
2. Pemetaan Pola Sebaran *Sand Dollar* dengan Menggunakan Citra Satelit Landsat di Pulau Menjangan Besar, Taman Nasional Karimun Jawa ..... 147
3. Kelimpahan dan Pola Sebaran *Echinodermata* di Pulau Karimunjawa, Jepara ..... 159
4. Struktur Komunitas Teripang (*Holothiroidea*) di Perairan Pulau Karimunjawa, Taman Nasioanl Karimunjawa, Jepara ..... 173

## Bencana Wilayah Pesisir, Laut dan Pulau-pulau Kecil: Ilmu Bencana dan Dampak Bencana

1. Kontribusi Nutrien N dan P dari Sungai Serang dan Wiso ke Perairan Jepara ..... 183
2. Kelimpahan, Keanekaragaman dan Tingkat Kerja Osmotik Larva Ikan pada Perairan Bervegetasi Lamun dan atau Rumput Laut di Perairan Pantai Jepara ..... 192
3. Pengaruh Fenomena Monsun, El Nino Southern Oscillation (ENSO) dan Indian Ocean Dipole (IOD) Terhadap Anomali Tinggi Muka Laut di Utara dan Selatan Pulau Jawa..... 205
4. Penilaian Pengkayaan Logam Timbal (Pb) dan Tingkat Kontaminasi Air Ballast di Perairan Tanjung Api-api, Sumatera Selatan ..... 218
5. KajianPotensi Energi Arus Laut di Selat Toyapakeh, Nusa Penida Bali ..... 225
6. Bioakumulasi Logam Berat Timpal pada Berbagai Ukuran Kerang *Corbicula javanica* di Sungai Maros ..... 235
7. Analisis Data Ekstrim Tinggi Gelombang di Perairan Utara Semarang Menggunakan *Generalized Pareto Disttribution* ..... 243
8. Kajian Karakteristik Arus Laut di Kepulauan Karimunjawa, Jepara ..... 254
9. Cu dan Pb dalam Ikan Juaro (*Pangasius polyuronodon*) dan Sembilang (*Paraplotosus albilabris*) yang Tertangkap di Sungai Musi Bagian Hilir, Sumatera Selatan..... 264
10. Kajian Perubahan Spasial Delta Wulan Demak dalam Pengelolaan Berkelanjutan Wilayah Pesisir..... 271
11. Biokonsentrasi Logam Plumbum (Pb) pada Berbagai Ukuran Panjang Cangkang Kerang Hijau (*Perna viridis*) dari Perairan Teluk Semarang..... 277



12. Hubungan Kandungan Bahan Organik Sedimen dengan Kelimpahan <i>Sand Dollar</i> di Pulau Cemara Kecil Karimunjawa, Jepara .....	287
13. Kandungan Logam Berat Kadmium (Cd) dalam Air, Sedimen, dan Jaringan Lunak Kerang Hijau ( <i>Perna viridis</i> ) di Perairan Sayung, Kabupaten Demak.....	301
<b>Bioteknologi Kelautan: Bioremediasi, Pangan, Obat-obatan .....</b>	
1. Pengaruh Lama Perendaman Kerang Hijau ( <i>Perna viridis</i> ) dalam Larutan Nanas ( <i>Ananas comosus</i> ) Terhadap Penurunan Kadar Logam Timbal (Pb) .....	312
2. Biodiesel dari Hasil Samping Industri Pengalengan dan Penepungan Ikan Lemuru di Muncar .....	328
3. Peningkatan Peran Wanita Pesisir pada Industri Garam Rebus .....	339
4. Pengaruh Konsentrasi Enzim Bromelin pada Kualitas Hidrolisat Protein Tinta Cumi-cumi ( <i>Loligo</i> sp.) Kering.....	344
5. Efek Enzim Fitase pada Pakan Buatan Terhadap Efisiensi Pemanfaatan Pakan Laju Pertumbuhan Relatif dan Kelulushidupan Ikan Mas ( <i>Cyprinus carpio</i> ).....	358
6. Substitusi Silase Tepung Bulu Ayam dalam Pakan Buatan Terhadap Laju Pertumbuhan Relatif, Pemanfaatan Pakan dan Kelulushidupan Benih Ikan Nila Larasati ( <i>Oreochromis niloticus</i> ) .....	372
7. Stabilitas Ekstrak Pigmen Lamun Laut ( <i>Enhalus acoroides</i> ) dari Perairan Teluk Awur Jepara Terhadap Suhu dan Lama Penyimpanan.....	384
8. Penggunaan Kitosan pada Tali Agel sebagai Bahan Alat Penangkapan Ikan Ramah Lingkungan .....	401
9. Kualitas Dendeng Asap Ikan Tongkol ( <i>Euthynnus</i> sp.), Tunul ( <i>Sphyrna</i> sp.) dan Lele ( <i>Clarias</i> sp.) dengan Metode Pengeringan Cabinet Dryer.....	408
<b>Aplikasi IPTEK Perikanan dan Kelautan dalam Pengelolaan dan Pemanfaatan Sumberdaya Wilayah Pesisir, Laut dan Pulau-pulau Kecil (Manajemen Sumberdaya Perairan)</b>	
1. Studi Karakteristik Sarang Semi Alami Terhadap Daya Tetas Telur Penyu Hijau ( <i>Chelonia mydas</i> ) di Pantai Paloh Kalimantan Barat .....	422
2. Struktur Komunitas Rumput Laut di Pantai Krakal Bagian Barat Gunung Kidul, Yogyakarta .....	434
3. Potensi dan Aspek Biologi Ikan Nila ( <i>Oreochromis niloticus</i> ) di Perairan Waduk Cacaban, Kabupaten Tegal.....	443



4. Morfometri Penyu yang Tertangkap secara <i>By Catch</i> di Perairan Paloh, Kabupaten Sambas, Kalimantan Barat.....	452
5. Identifikasi Kawasan <i>Upwelling</i> Berdasarkan Variabilitas Klorofil-A, Suhu Permukaan Laut dan Angin Tahun 2003 – 2015 (Studi Kasus: Perairan Nusa Tenggara Timur).....	463
6. Hubungan Kelimpahan Fitoplankton dan Zooplankton di Perairan Pesisir Yapen Timur Kabupaten Kepulauan Yapen, Papua.....	482
7. Analisis Hubungan Kandungan Bahan Organik dengan Kelimpahan Gastropoda di Pantai Nongsa, Batam .....	495
8. Studi Morfometri Ikan Hiu Tikusan ( <i>Alopias pelagicus</i> Nakamura, 1935) Berdasarkan Hasil Tangkapan di Pelabuhan Perikanan Samudera Cilacap, Jawa Tengah.....	503
9. Variabilitas Parameter Lingkungan (Suhu, Nutrien, Klorofil-A, TSS) di Perairan Teluk Tolo, Sulawesi Tengah saat Musim Timur.....	515
10. Keanekaragaman Sumberdaya Teripang di Perairan Pulau Nyamuk Kepulauan Karimunjawa .....	529
11. Keanekaragaman Parasit pada Kerang Hijau ( <i>Perna viridis</i> ) di Perairan PPP Morodemak, Kabupaten Demak .....	536
12. Model Pengelolaan Wilayah Pesisir Berbasis Ekoregion di Kabupaten Pemalang Provinsi Jawa Tengah .....	547
13. Ektoparasit Kepiting Bakau ( <i>Scylla serrata</i> ) dari Perairan Desa Wonosari, Kabupten Kendal.....	554
14. Analisis Sebaran Suhu Permukaan Laut, Klorofil-A dan Angin Terhadap Fenomena <i>Upwelling</i> di perairan Pulau Buru dan Seram...	566
15. Pengaruh Pergerakan Zona Konvergen di Equatorial Pasifik Barat Terhadap Jumlah Tangkapan Skipjack Tuna ( <i>Katsuwonus pelamis</i> ) Perairan Utara Papua – Maluku.....	584
16. Pemetaan Kandungan Nitrat dan Fosfat pada Polip Karang di Kepulauan Karimunjawa .....	594
17. Hubungan Kandungan Bahan Organik dengan Distribusi dan Keanekaragaman Gastropoda pada Ekosistem Mangrove di Desa Pasar Banggi Kabupaten Rembang.....	601

#### Aplikasi IPTEK Perikanan dan Kelautan dalam Pengelolaan dan Pemanfaatan Sumberdaya Wilayah Pesisir, Laut dan Pulau-pulau Kecil (Budidaya Perairan)

1. Pengaruh Suplementasi <i>Lactobacillus</i> sp. pada Pakan Buatan Terhadap Aktivitas Enzim Pencernaan Larva Ikan Bandeng ( <i>Chanos chanos</i> Forskal).....	611
2. Inovasi Budidaya Polikultur Udang Windu ( <i>Penaeus monodon</i> ) dan Ikan Koi ( <i>Cyprinus carpio</i> ) di Desa Bangsri, Kabupaten Brebes: Tantangan dan Alternatif Solusi.....	621





3. Pertumbuhan dan Kebiasaan Makan Gelondongan Bandeng ( <i>Chanos chanos</i> Forskal) Selama Proses Kultivasi di Tambak Bandeng Desa Wonorejo Kabupaten Kendal .....	630
4. Analisis Faktor Risiko yang Mempengaruhi Serangan <i>Infectious Myonecrosis Virus</i> (IMNV) pada Budidaya Udang Vannamei ( <i>Litopenaeus vannamei</i> ) secara Intensif di Kabupaten Kendal .....	640
5. Respon Histo-Biologis Pakan PST Terhadap Pencernaan dan Otak Ikan Kerapu Hibrid ( <i>Epinephelus fuscoguttatus</i> x <i>Epinephelus polyphekaidon</i> ).....	650
6. Pengaruh Pemberian Pakan <i>Daphnia</i> sp. Hasil Kultur Massal Menggunakan Limbah Organik Terfermentasi untuk Pertumbuhan dan Kelulushidupan ikan Koi ( <i>Carassius auratus</i> ) .....	658
7. Pengaruh Aplikasi Pupuk NPK dengan Dosis Berbeda Terhadap Pertumbuhan <i>Gracilaria</i> sp. ....	668
8. Pengaruh Vitamin C dan <i>Highly Unsaturated Fatty Acids</i> (HUFA) dalam Pakan Buatan Terhadap Tingkat Konsumsi Pakan dan Pertumbuhan Ikan Patin ( <i>Pangasius hypophthalmus</i> ) .....	677
9. Pengaruh Perbedaan Salinitas Media Kultur Terhadap Performa Pertumbuhan <i>Oithona</i> sp. ....	690
10. Mitigasi Sedimentasi Saluran Pertambakan Ikan dan Udang dengan Sedimen Emulsifier di Wilayah Kecamatan Margoyoso, Pati .....	700
11. Performa Pertumbuhan <i>Oithona</i> sp. pada Kultur Massal dengan Pemberian Kombinasi Pakan Sel Fitoplankton dan Organik yang Difermentasi.....	706
12. Respon Osmotik dan Pertumbuhan Juvenil Abalon <i>Haliotis asinina</i> pada Salinitas Media Berbeda.....	716
13. Pengaruh Pemuasaan yang Berbeda Terhadap Pertumbuhan dan Kelulushidupan Ikan Nila ( <i>Oreochromis niloticus</i> ) .....	728

**Aplikasi IPTEK Perikanan dan  
Kelautan dalam Pengelolaan dan  
Pemanfaatan Sumberdaya Wilayah  
Pesisir, Laut dan Pulau-pulau Kecil  
(Pemanfaatan Sumberdaya Perairan)**



## EFFECT OF DIFFERENT SOAKING TIME IN COCONUT SHELL LIQUID SMOKE TO THE PROFILE OF LIPIDS CAT FISH (*CLARIAS BATRACHUS*) SMOKE

Sarah Nur Halimah\*)<sup>1</sup>, Fronthea Swastawati<sup>1</sup>, Romadhon<sup>1</sup>

<sup>1</sup>Departmen of Fisheries Diponegoro University, Jl.Prof Soedarto, SH Semarang-50275, Indonesia

\* Correspondence : Phone +62-85-640590491; e-mail : sarahnurhalimah@gmail.com

### ABSTRACT

Catfish is a freshwater fish that contain fat quite high due to the feed. Catfish fat can affect oxidation, end product quality smoked of fish. The addition of liquid smoke in the smoking process was able to maintain the fatty acids, especially unsaturated fatty acids. The compounds in the liquid smoke serves to inhibit and prevent the oxidation of fat. Phenol content in liquid smoke works by stabilizing free radicals, thus preventing damage to fat. The material used in this study are catfish, coconut shell liquid smoke, salt, and water. The experimental design used is Complete Random Design (RAL), which consists of one treatment (long immersion in liquid smoke) 0 hours, 1 hour, 2 hours, and 3 hours with repeat 3 times. Parameter testing include test fat levels, fatty acids profile, cholesterol, phenol content, water content, and hedonic. The results showed that long immersion in different liquid smoke significant effect on levels of fat, cholesterol, phenol levels, and fatty acids profile. The value of the fat content of each treatment 0 hour, 1 hour, 2 hours, and 3 hours were  $5.39 \pm 0.11\%$ ;  $6.76 \pm 0.08\%$ ;  $9.32 \pm 0.37\%$ ;  $11.43 \pm 0.11\%$ . Values saturated fatty acid profile of 29.27%, 37.61%, 38, 90%, and 39.20%. Unsaturated fatty acids of 38.16%, 42.53%, 52.10%, and 54.13%. Cholesterol values of  $13,5 \pm 0,26\text{mg}/100$ ;  $10,35 \pm 0,13\text{mg}/100$ ;  $8,12 \pm 0,10\text{mg}/100$ ;  $5,15 \pm 0,93\text{mg}/100$ . Water content of  $62.09 \pm 0.90\%$ ;  $56.44 \pm 2.28\%$ ;  $52.40 \pm 0.57\%$ ;  $51.99 \pm 1.08\%$ . Value phenol content of  $2,18 \pm 0,05\text{mg}/\text{l}$ ;  $3,39 \pm 0,14\text{mg}/\text{l}$ ;  $5,33 \pm 0,37\text{mg}/\text{l}$ ;  $6,47 \pm 0,32\text{mg}/\text{l}$ . Fish treatment is most effective for maintaining the quality of fatty acids in general are soaking time 2 hours

**Keywords: Catfish, Coconut Shell Liquid Smoke, Fatty Acid Profile**

### I. INTRODUCTION

Fishing is an important sector in foreign exchange earnings to the development of Indonesia. Fish has an important role in the improvement of human nutrition. The production of aquaculture in 2009 reached 4.71 million tons. Lele fish farming in Central Java is increasing every year, so that the processed product and catfish fish consumption is also higher. In 2010, production Catfish in Central Java province from aquaculture pond reaches 36394.5 tons. Demak is one of the most highly rated producers farming catfish fish production amounted to 12,550 tons (Directorate General of Aquaculture, 2013).

Catfish is one of fishery products that contain fatty acids that are quite high. Fatty acids are long-chain organic acids having a carboxyl group (COOH) at one end and a methyl group (CH<sub>3</sub>) at the other end. Processed fish catfish very loved by the people because the price of fish catfish is quite affordable and has a savory taste, so it has been used optimally into smoked fish and processed food diversification. According Murniyati *et al.* (2013), catfish fat can lower LDL (Low Density Lipids) cholesterol in the blood



plasma. Fish fatty acids generally straight carbon chain and have carbon atoms between 14-24 atoms.

The fish is needed by the human body because it has some benefits, among others educate the brain, helps the growth period, and lower triglyceride levels. One of the processed fish to maintain the content of fatty acids is a smoked fish. Curing fish with liquid smoke can inhibit the oxidation of fat because it contains phenols, organic acids, and antioxidants, so that evaporation can maintain-acid content of unsaturated fatty acids required for the body. According Darmadji (1999), liquid smoke coconut shell proven to have a major distinction in terms of the intensity of the color, smell, and taste-specific and was followed by the ability to inhibit the growth of mold and fat oxidation.

## II. MATERIALS AND METHODS

### 2.1. Smoking process

The fish were dipped in the 5 % brine solution, and then dipped in the 1 % liquid smoke solution. Liquid smoke was produced from coconut shell. The dipping process were divided into three groups different soaking time: (i) the fish dipped in liquid smoke without soaking shell as control; (ii) the fish dipped in liquid smoke soaking time in one hours (1h); (iii) the the fish dipped in liquid smoke soaking time in two hours (2h); and (iv) the the fish dipped in liquid smoke soaking time three hours (3h). The processing time and temperature in the kiln were at 90 °C (3 h). After heating and cooling process, the fatty acid and quality (lipid; water; phenol and hedonic content) of smoked fish were observed.

### 2.2. Fatty acid composition

Approximately 25 mg of fat, obtained from the lipid extraction according to Folch *et al.* (1957), were saponified, and the fatty acids converted to methyl esters according to Joseph and Ackman (1992). The methyl esters were dissolved in hexane and 2 mL were injected into a gas chromatograph with a flame ionization detector (GC-FID) (Shimadzu, model GC 2010, Kyoto, Japan). The methyl esters were separated in a capillary column of melted silica (100 m \_ 0.25 mm i.d., 0.20 mm thickness of the stationary phase) (CP-SIL 88, Chromopack, Middleburg, The Netherlands) and identified by comparison of the retention times of the sample peaks with those of standards (FAME Mix C4eC24, Supelco, Bellefonte, Pennsylvania, USA). The quantification was carried out by internal standardization using the tridecanoic acid methyl ester as internal standard added before sample injection. The results were calculated in mg per 100 g of sample (AOCS, 1989).





### 2.3. Chemical composition

#### 2.3.1. Total Lipid Content

Total Lipid Content was determined by Soxhlet extraction method using petroleum ether solvent (Soxhlet, 1879) and by Bligh and Dyer method using methanol-chloroform (2:1, v/v) (Bligh & Dyer, 1959).

#### 2.3.2. Cholesterol

Cholesterol content was estimated using Liebermann-Burchard reagent (Attarde *et al.*, 2010). Standard cholesterol solution used was 2 mg/ml as stock solution. Liebermann-Burchard reagent was prepared with 7 ml concentrated sulfuric acid and 5 ml glacial acetic acid and was covered with black paper and kept in ice bucket in dark place.

#### 2.3.3. Water content

Determination of water smoked fish were carried out in triplicate according to AOAC 2005. Water content of the samples were determined by method in which 2 g of the samples (fish muscles) were oven-dried at  $110 \pm 1^\circ\text{C}$  for 3 h to a constant weight. Loss in weight is equal to the water content of the original sample.

#### 2.3.4. Total phenol

The total phenolic content were determined by using the Folin Ciocalteu assay. A reagent blank using distilled water was prepared. 1 ml of Folin-Ciocalteu phenol reagent was added to the mixture and shaken. After 5 minutes 10 ml of 7%  $\text{Na}_2\text{CO}_3$  solution was added to the mixture. The volume was then made up to the mark. After incubation for 90 minutes at room temperature, the absorbance against the reagent blank was determined at 765 nm with an UV-Visible spectrophotometer.

### 2.4. Hedonic test

The organoleptic evaluation of the smoked samples was carried out using a 30 member panelists. Parameters such as flavour, texture, odour, appearance and general taste were used to compare the organoleptic characteristics of the products. Questionnaires were used by the panelists and scoring was done on a weekly basis. The questionnaires were prepared using a 5 point hedonic scale as suggested by Poste *et al.* (1991). The points are as outlined below: Very good – 5, Good – 4, Averagely good – 3, Fair – 2, and Bad – 1

### 2.5 Statistical Analysis

Experimental fishes were laid out in a completely randomized design. The data obtained were subjected to analysis of variance (ANOVA) and using statistical package for social science (SPSS) Version 16.0 computer software.



### III. RESULT

The preliminary study was conducted to determine the best concentration of Catfish smoked with concentration (1%); (3%); and (5%) by carrying out organoleptic tests include appearance, odor, flavor, texture, fungi, slime and concluded that the highest value and the best of the three treatments is the concentration of 1% to be used for primary research. The main research uses a 1% concentration with a long treatment difference immersion in liquid smoke that is control, 1 hour, 2 hours, and 3 hours in cold temperatures.

#### 3.1. Fatty Acid Analysis

The results indicate that the fatty acid profile obtained has two types of fatty acids. Class of fatty acids are found consisting of saturated fatty acids and unsaturated fatty acids. Difference between the two lies in their chemical bonds, where the saturated fatty acid has no double bonds. These differences led to their physical and chemical properties of both these fatty acids. Based on the results of Table 1 shows the saturated fatty acids of fish catfish on difference in the treatment of soaking liquid smoke for control, 1 hour, 2 hours, and 3 hours on average by 29.27%, 37.61%, 38, 90%, and 39 , 20%, while unsaturated fatty acids are as follows: control, 1 hour, 2 hours, and 3 hours on average by 38.16%, 42.53%, 52.10% and 54.13%. Differences in the composition of fatty acids in each sample was highly dependent on the source of these fats. According Pujiati *et al.* (2005), the composition of fatty acids remain in the fish that have been preserved using liquid smoke. The smoked fish was still able to meet essential fatty acid needs. Palmitoleic acid, 11-oktadekenoat acid, and oleic acid is an unsaturated fatty acid. Palmitic acid, isostearic acid and stearic acid is a saturated fatty acid. Arachidonic acid (ARA) and Eikosapetaenoat acid (EPA) is an omega-3 fatty acids, essential fatty acids whose role is very important for the body, among others, are useful in the development of the brain and nervous sense of sight.

**Tabel 1. Fatty Acid Catfish Smoke**

Fatty Acid	Fatty acid content (%) in the immersion liquid smoke			
	control	1 hour	2 hours	3 hours
<b>1.Saturated Fatty Acids</b>				
Myristic acid (C14:0)	1,51	1,25	2,18	1,65
Palmitate acid (C16:0)	18,75	24,86	28,16	29,22
Stearat acid (C18:0)	6,95	7,87	7,78	6,56
Laurat acid (C12:0)	2,06	3,63	0,78	1,77
<b>Jumlah</b>	29,27	37,61	38,90	39,20
<b>2. Unsaturated Fatty Acids</b>				



**- Mono Unsaturated Fatty Acids**


Elaidic acid (C18:1n-9)	22,30	24,77	26,12	32,27
Palmitoleic acid (C16:1n-7)	2,06	1,45	3,19	1,36
Oleat acid (C18:1n-9)	0,78	3,09	3,83	5,42

**-Poly Unsaturated Fatty Acids**

*Linoleat acid (C18:2n-6)	10,59	11,18	11,98	11,47
*arachidonic acid (C20:4n-6)	1,66	0,67	2,84	0,64
*EPA (C20:5n-3)	0,77	1,37	4,14	2,97

<b>Jumlah</b>	<b>38,16</b>	<b>42,53</b>	<b>52,10</b>	<b>54,13</b>
---------------	--------------	--------------	--------------	--------------

Description: \* sign an essential fatty acid



**Figure 1.** Fatty Acid Profile

## 1. Saturated Fatty Acid

The content of saturated fatty acids in each treatment shows the results of different it is because these fatty acids are the basic components of the formation of fat in fish catfish, while the content of unsaturated fatty acids in fish are high can be caused due to the mechanism of the food chain and the activity of the fish high, causing large enough fat reserves. According Afrianto and Liviawaty (2005), need fat in fish is influenced by environmental conditions, feed and seasonal variations. Differences fatty acids in marine and freshwater fish related to the specific needs of these fish to adapt physiologically to the environment. Feed also affect the composition of fatty acids in fish.

Results showed that saturated fatty acid palmitic acid has a percentage value compared with other saturated fatty acids. Palmitic fatty acid is a long chain saturated fatty acids contained in animal fats. Foods that contain lots of saturated fat is not good for health. According Jacob *et al.* (2014), shows the palmitic acid is a saturated fatty acid (SFA) with

the highest levels, both in the flesh of fresh eel and eel stew. Fresh eel contains palmitate 13.79% and subject to change after the boiling process becomes 12.88%. Palmitic acid is a saturated fatty acid (SFA) is most commonly found in foodstuffs, ie (15-50)% of total fatty acids that exist (Winarno 2008).

## 2. Unsaturated Fatty Acid

Unsaturated fatty acids are fatty acids that have the double bond in their chemical structure. Unsaturated fatty acids are divided into two types, namely monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA). MUFA is an unsaturated fatty acid having only one double bond, whereas PUFAs are fatty acids that have more than one double bond. PUFAs made up of fatty acids omega-3 (n-3), omega-6 (n-6), and omega-9 (N-9), a growing number of double bonds in unsaturated fat, the more vulnerable nature against fat oxidation, in addition to the unsaturated fatty acids will also be damaged if exposed to heat. By Arpi (2014), oxidation easily occur in the processing, storage, and use of oil in food fats, oxidation will cause rancidity. Prevention of oxidation of fatty food can be done with good handling and storage, among others at the appropriate temperature, not exposed to light, low water content, and the absence of a metal catalyst. Oxidation can also be prevented by the addition of antioxidants, which is a compound in a small amount can slow oxidation in the material. Certain food ingredients contain antioxidants naturally, but natural antioxidant content generally decreased during the treatment process. Eicosapentaenoic acid (EPA) adalah dua rantai panjang asam lemak tak jenuh ganda milik Omega-3.

EPA content is determined by the type of fatty fish used. Table 1 shows that the EPA in the treatment of soaking liquid smoke on the control to 3 hours soaking time is getting higher. Control by 0.77%, 1 hour by 1.37%, 2.84% at 2 hours and 3 hours by 2.97%, but the soaking time 2 hours to 3 hours with smoked catfish fish began to decline. This is because the longer the oxidation of fish catfish decomposed due to the fat composition is different. Unsaturated fatty acids lead to fats in fish is easily oxidized. The oxidation process can cause the flavor and taste are not preferred as well as a decrease in nutritional value. This is consistent with research Purwaningsih *et al.* (2014), the effect of fish processing affect the content of unsaturated fatty acids such as fish Glodohk, EPA fresh fish 11.14%, 12.46% salt boiled, cooked without treatment 12.08%, 13.50% and steamed. According Kolalowska (2003), fat oxidation stability is influenced by internal and external factors, such as fatty acid composition, content of prooxidant, antioxidant,





irradiation, temperature, oxygen, the surface area in contact with oxygen, the level of processing and storage conditions.

### 3.2. Chemical Composition Analysis

**Table 2. Test Results fish Catfish Smoke**

Test		Treatment			
		control	1 hour	2 hours	3 hours
Total Lipid	Content (%)	5,39±0,11a	6,76±0,08b	9,32±0,37c	11,43±0,11d
Cholesterol	(mg/10g)	13,5±0,26a	10,35±0,13b	8,12±0,10c	5,15±0,93d
Water content	(%)	62,09±0,90a	56,44±2,28b	52,40±0,57c	51.99±1,08c
Total phenol	(mg/l)	2,18±0,05a	3,39±0,14b	5,33±0,37c	6,47±0,32d

Information:

- Data is the average of three replicates  $\pm$  standard deviation
- Data followed different letters indicate significant differences ( $\leq 0.05$ )

#### *Analysis of Fat Content*

The content of the highest fat content contained in the long immersion liquid smoke 3 hours 11.43% and the lowest is the long soaking liquid smoke for 0 hours. The influence of the value of the fat content in fish smoked catfish in each treatment due to their phenolic compounds seep into the structure of the fish meat catfish. Aktivitas phenol having antioxidant that can prevent the oxidation reaction. The oxidation reaction is influenced by phenol antioxidant compounds that provide hydrogen groups in preventing fat oxidation and damage. Small big influence results of the fat content is determined by the water content contained in the fish, the evaporation process can generally reduce the water content of the composition as a result of the heat treatment process. The heat that comes from the fuel will penetrate into the network structure of fish meat, then the water will come out. According Yusnaini *et al.* (2015), the water content in the process of curing fish smaller than that of the fresh material due to the method of heating and the final temperature that is used.

#### *Analysis of Cholesterol Levels*

Cholesterol levels were obtained by long immersion liquid smoke for 1 hour, 2 hours, and 3 hours respectively is 10.35 mg / 100g, 8.12 mg / 100g, and 5.15 mg / 100g. Lele fish cholesterol content analysis was conducted to determine the treatment prolonged submersion different liquid smoke can lower cholesterol levels in fish catfish. Based on the results of Table 10 decrease in cholesterol levels were highest occurred in the commission of 3 hours soaking time is equal to 5.15 mg / 100g, while the percentage decrease in



cholesterol levels was lowest in controls (0 hour soaking time), so it was concluded that the cholesterol content than control experience real difference. Decrease in cholesterol levels caused by the content of phenol-containing antioxidant. This antioxidant substances could inhibit oxidation reactions by binding free radicals and reactive molecules. According Swastawati (2009), a decrease in cholesterol content can be due administration of liquid smoke in the smoking process, because the phenolic compounds contained in the liquid smoke can break the chains of fat oxidation in the initiation phase. Phenolic compounds contained in liquid smoke are antioxidants and antibacterial.

#### *Results Analysis of Water Content*

The results of water content in fish catfish showed that the water content control (0 hours) has exceeded the limits of the standard Indonesian National Standard (2013) is a maximum of 60%. Results of treatment of soaking liquid smoke at 0% has a value of 62.09%, while the treatment of 1 hour, 2 hours and 3 hours is 56.44%, 52.40% and 51.99%. This is in comparison research Syarafina *et al.* (2014), Tengiri smoked fish with a value of liquid smoke soaking time of 5 minutes 50.85%, 48.72% 10 minutes, and 15 minutes 47.62%. According Swastawati (2011), stated that the decrease in water levels also dipengaruhi by their treatment prior to curing such as draining and soaking in the concentration of certain salts

#### *Results Analysis of Phenol Content*


The results of the analysis show that the phenol content of phenol levels with long soaking liquid smoke different on catfish smoke rose. This is due to the longer time soaking smoked catfish fish the more phenol compounds are absorbed into the catfish. Phenol compounds are in liquid smoke coconut tempuung are antioxidants that inhibit the oxidation of fatty fish catfish processed fish products cause damage. Phenol was instrumental in forming the taste and aroma of fisheries products. According Soazo *et al.* (2015), a liquid smoke product comprising phenol, carbonyl component and acetic acid. Phenol functions as an antioxidant and antibacterial food. Lira *et al.* (2013), stating the amount of salt, hot distance, time of fumigation will affect the curing phenol but no effect on the increase in cholesterol, the cholesterol is determined by polyunsaturated fatty acids only found on shrimp.

The results showed that the effectiveness of phenol greatest lies in the long immersion liquid smoke 2 hours. Value effectiveness of phenol can be seen on the biggest difference on a long hose soaking liquid smoke that has a difference of 1,94. Based on the results table is known that phenol levels in 3 hours soaking time of 6,47mg/l is the largest



phenol content. According Swastawati et al. (2009), vinegar has a high phenol content does not always produce smoked fish with high levels of phenols. Instead of smoke that have a low phenol content can be absorbed more into the flesh of fish depend on the nature and texture of the flesh of fish biology, in the application of fumigation concentration used is 1-3%, which is sufficient for consumers.

### 3.3. Hedonic Test



**Figure 2.** Diagram Hedonic Test

**Table 3.** Data Value Hedonic Test Catfish Smoke Based Differences in Old Soaking Liquid Smoke

Specification	Soaking Liquid Smoke			
	Control	1 hour	2 hours	3 hours
Appearance	5,90±0,75a	6,86±0,62b	7,26±0,63c	8,10±0,71d
Flavor	5,73±0,83a	6,96±0,61b	7,46±0,73bc	8,06±0,69d
Smell	5,63±0,61a	7,03±0,67b	7,30±0,70bc	7,97±0,76d
Texture	5,86±0,77a	6,90±0,61b	7,20±0,66bc	7,93±0,63d
Average	5,78±0,74	6,93±0,62	7,31±0,68	8,02±0,70

Based on the hedonic test results against control smoked catfish fish at test level obtained 95% confidence interval of  $5.648 \leq \mu \leq 5.918$ , so it can be concluded that fish smoked catfish are less preferred by the panelists. Soaking time 1 hour on the obtained confidence interval of  $6.851 \leq \mu \leq 7.033$ , the soaking time 2 hours gained the confidence interval of  $7.170 \leq \mu \leq 7.446$ , and the soaking time 3 hours gained the confidence interval of  $7.862 \leq \mu \leq 8.172$ , so it concluded that the treatment 3 hours soaking time provide results that most preferred by the panelists.

The resulting texture in smoked catfish fish that is dense, compact, fairly elastic, and not loud. This texture is affected due to long curing time is causing water levels of smoke is reduced, causing a mushy texture. Cai long immersion in the smoke will affect the amount of phenol produced, the longer the curing time will cause phenol components that attach to the fish will be many more. According Adiyastiti *et al.* (2014), the physical properties of the meat, such as color, texture, hardness and tenderness of meat is affected by the binding power of water, the water content of the matrix protein's ability to hold

water or absorb water being added from outside influences such as cooking. Violence is the deciding factor in the formation of the texture of a product.

## Conclusion

Giving long difference in liquid smoke terhadap Catfish (*Clarias batrachus*) Smoke significant effect on the decline in fish fat profile catfish. The use of liquid smoke time difference is 1 hour, 2 hours, and 3 hours tend to lower cholesterol levels and water content, maintaining the levels of fat, as well as maintaining the unsaturated fatty acids and EPA on smoked catfish.

## References

- Adiyastiti, B.E.T., E. Suryanto, dan Rusman. 2014. Pengaruh Lama Pembakaran dan Jenis Bahan Bakar terhadap Kualitas Sensoris dan Kadar Benzo (A) Piren Sate Daging Kambing. *Buletin Peternakan*, 33(3): 189-196 ISSN: 0126-4400.
- Afrianto, E. dan E. Liviawaty. 2010. *Penanganan Ikan Segar*. Widya Pajaran, Bandung, 156 hlm. ISBN: 978-602-8323-55-0.
- Arpi, N. 2014. Kombinasi Antioksidan Alamai a-Tokoferol dengan Asam Askorbat dan Antioksidan Sintesis BHA dalam Menghambat Ketengikan Kelapa Gongseng Giling (U Neulheu) Selama Penyimpanan. *Jurnal Teknologi dan Industri Pertanian Indonesia*, Universitas Syiah Kuala, 6 (2).
- Association of Official Analytical Chemist [AOAC]. 2005. *Official Method of Analysis of The Association of Official Analytical of Chemist*. Arlington: The Association of Official Analytical Chemist, Inc.
- Attarde MA. 2010. *Physiological Effects of Some Plant Oils Supplementation on Streptozotocin-Induced Diabetic Rats*. *Res. J. Med. Sci.*, 5: 55-71.
- Badan Standardisasi Nasional. 2013. SNI 2725-2013. Ikan Asap dengan Pengasapan Panas. Syarat Mutu dan Keamanan Produk. Badan Standardisasi Nasional. Jakarta.
- Bligh, E. G., and Dyer, W. J. 1959. *A Rapid Method of Total Lipid Extraction and Purification*. *Canadian Journal of Biochemistry Physiology*, 37, 911-991. <http://dx.doi.org/10.1139/o59-099>
- Darmadji, P., 1999. Sifat Antioksidatif Asap Cair Hasil Redistilasi Selama Penyimpanan. Prosiding Seminar Nasional Pangan, Pusat Antar Universitas Pangan dan Gizi. UGM. Yogyakarta.
- Direktorat Jendral Perikanan dan Kelautan. 2013. Profil Kelautan dan Perikanan Provinsi Jawa Tengah Untuk Mendukung Industrialisasi KP. Jakarta.
- Folch, J., Lees, M., and Stanley, S. A. 1957. *Simple Method for The Isolation and Purification of Total Lipide from Animal Tissues*. *The Journal of Biological Chemistry*.
- Jacob, A.M., P. Suptijah, dan R. Kamila. 2014. Kandungan Asam Lemak, Kolesterol, dan Deskripsi Jaringan Daging Belut Segar dan Rebus. *JPHPI*, 17 (2): 134-143.
- Joseph, J. D., and Ackman, R. G. 1992. *Capillary Column Gas Chromatographic Method for Analysis of Encapsulated Fish Oils and Fish Oil Ethyl Esters: Collaborative Study*. *Journal of AOAC International*, 75.
- Kolalowska A. 2003. *Lipid Oxidation In Food Systems*. CRC Press. Washington DC.





- Lira, G.M., K.W.B. Silva, B.C. Figueiredo, and N. Bragagnolo. 2014. *Impact of smoking on the lipid fraction and nutritional value of seabob shrimp (Xiphopenaeus kroyeri, Heller, 1862)*. *Journal Food Science and Technology*. Elsevier. 183-187.
- Murniyati, Th. D. Suryaningrum, dan I. Muljanah. 2013. *Membuat Fillet Lele dan Produk Olahannya*. Penebar Swadaya, Jakarta, 84 hlm. ISBN 979-002-637-4.
- Poste LM, Mackie DA, and Butler G. 1991. *Laboratory Methods for Sensory Analysis of Food*. Ottawa. Res. Branch Agric., Canada Publication.
- Pujiati, P. P.R. Istianingrum, dan A.A. Irawan. 2005. Pengaruh Pengawetan Ikan Kembung (*Rastrelliger Neglectus*) dengan Asap Cair Tempurung Kelapa Terhadap Penghambatan Kerusakan Protein, Kadar Lemak dan Komposisi Asam Lemaknya. *Jurnal PELITA*. 1. (1). Universitas Negeri Yogyakarta.
- Purwaningsih, S., E. Salamah, dan R. Dewantoro. 2014. *Komposisi Kimia dan Asam Lemak Ikan Glodok Akibat Pengolahan Suhu Tinggi*. JPHPI. Institut Pertanian Bogor. 17 (2).
- Soazo, M. L.M Perez, G.N. Piccirilli, N.J. Deloenzi, and R.A. Verdini. 2015. *Antimicrobial and physicochemical characterization of whey protein concentrante edible films incorporated with liquid smoke*. *Journal LWT-Food Science and Technology*. Elsevier. 285-291.
- Swastawati., Y.S. Darmanto, T.W. Agustini, dan E. N. Dewi. 2009. *Pengasapan Ikan dari Tradisional sampai Modern*. Badan Penerbit UNDIP, Semarang, 140 hlm. ISBN: 978-979-704-871-6.
- \_\_\_\_\_, F. 2011. Studi Kelayakan Dan Efisiensi Usaha Pengasapan Ikan dengan Asap Cair Limbah Pertanian. *Jurnal Dinamika Ekonomi Pembangunan*, 1(1). Hal: 18-24.
- Syarafina, I. L., F. Swastawati, dan Romadhon. 2014. Pengaruh Daya Serap dan Lama Perendaman Yang Berbeda Terhadap Kualitas Dendeng Ikan Bandeng (*Chanos chanos* Forsk) dan Ikan Tengiri (*Scomberomorus* sp) Asap. *Jurnal Pengolahan dan Bioteknologi Hasil Perikanan*, 3 (1): 50-59.
- Winarno F.G. 2008. *Kimia Pangan dan Gizi*. Gramedia Pustaka Utama, Jakarta.
- Yusnaini, Soeparno, E. Suryanto, and R. Armunanto. 2014. *The Effect of Heating Poces Using Electric and Gas Ovens On Chemical and Physical Poperties of Cooked Smoked-Meat*. *Journal Procedia Food Science*. Elsiwier. 19-26.



